

SECRET

DRT DOCUMENTATION PAGE

1a. AD-A223 217			1b. RESTRICTIVE MARKINGS	
2a. SELECTED			3. DISTRIBUTION/AVAILABILITY OF REPORT	
2b. DECLASSIFICATION/DOWNGRADING SCHEDULE JUN 15 1990			Approved for public release; distribution unlimited.	
4. PERFORMING ORGANIZATION REPORT NUMBER(S)			5. MONITORING ORGANIZATION REPORT NUMBER(S)	
6a. NAME OF PERFORMING ORGANIZATION University of Minnesota			7a. NAME OF MONITORING ORGANIZATION U. S. Army Research Office	
6b. OFFICE SYMBOL (If applicable)			7b. ADDRESS (City, State, and ZIP Code) P. O. Box 12211 Research Triangle Park, NC 27709-2211	
6c. ADDRESS (City, State, and ZIP Code) Dept. of Chemical Eng. & Materials Sci. Minneapolis, MN 55455			9. PROCUREMENT INSTRUMENT IDENTIFICATION NUMBER DAAL03-86-G-0034	
8a. NAME OF FUNDING/SPONSORING ORGANIZATION U. S. Army Research Office			10. SOURCE OF FUNDING NUMBERS	
8b. OFFICE SYMBOL (If applicable)			PROGRAM ELEMENT NO.	
8c. ADDRESS (City, State, and ZIP Code) P. O. Box 12211 Research Triangle Park, NC 27709-2211			PROJECT NO.	
11. TITLE (Include Security Classification) Diffusion of Penetrants in Polymers			TASK NO.	
12. PERSONAL AUTHOR(S) Tirrell, Matthew, Frick, Theodore S. and Lodge, Timothy P.			WORK UNIT ACCESSION NO.	
13a. TYPE OF REPORT FINAL			14. DATE OF REPORT (Year, Month, Day) 1990 May 7	
13b. TIME COVERED FROM 2/3/86 TO 2/1/90			15. PAGE COUNT 2	
16. SUPPLEMENTARY NOTATION The view, opinions and/or findings contained in this report are those of the author(s) and should not be construed as an official Department of the Army position, policy, or decision, unless so designated by other documentation.				
17. COSATI CODES			18. SUBJECT TERMS (Continue on reverse if necessary and identify by block number)	
FIELD	GROUP	SUB-GROUP	glassy polymers, self-diffusion, probe dye molecule, Vrentas-Duda free volume theory, Penetrant Materials, Holographic Relaxation of Polymers, Polystyrene	
19. ABSTRACT (Continue on reverse if necessary and identify by block number)				
<p>Much effort has been expended in trying to understand and predict the unusual phenomena which can occur when initially glassy polymers are exposed to small-molecule gases, organic vapors or liquids. Sorption of the gas or vapor seems to occur at two distinctly different sorption sites (dual mode sorption), and influx of the penetrant leads to a non-Fickian front traveling at constant velocity (case-II diffusion). Because traditional measurement techniques such as sorption and permeation are complicated by these very phenomena which they try to explain, attempts to determine to what extent these phenomena are governed by diffusive characteristics of glassy transport rather than by the interplay of diffusion and polymer relaxation have been frustrated. Instead, a method of observing diffusion which does not require penetrant transport across a surface and influx into the sample must be used.</p> <p>(continued on reverse)</p>				
20. DISTRIBUTION/AVAILABILITY OF ABSTRACT <input type="checkbox"/> UNCLASSIFIED/UNLIMITED <input type="checkbox"/> SAME AS RPT. <input type="checkbox"/> DTIC USERS			21. ABSTRACT SECURITY CLASSIFICATION Unclassified	
22a. NAME OF RESPONSIBLE INDIVIDUAL			22b. TELEPHONE (Include Area Code)	
			22c. OFFICE SYMBOL	

(Block 19 continued)

We have used the Holographic Relaxation Spectroscopy (HRS) technique to eliminate the stumbling blocks mentioned above. HRS allows us to observe the self-diffusion of a probe dye molecule dispersed homogeneously in a polymer or polymer-diluent mixture at equilibrium, thus eliminating complications mentioned above. The diffusion is measured on length scales in the micron range, by creating a spatially periodic variation in dye concentration with a flash of laser light; the rate of disappearance of this pattern is proportional to the diffusion coefficient. Using HRS, we can measure self-diffusion coefficients (D_{self}) ranging from 10^{-5} to 10^{-16} cm²/sec. This allows measurement in systems ranging from pure diluent to glassy bulk polymer. With the capabilities of HRS at our disposal, we are addressing the following questions concerning the nature of small molecule transport in glassy and non-glassy systems:

- 1) What is the effect of diffusant molecular size on its transport?
- 2) What is the effect of the glass transition on $D_{\text{self}}(c, T)$?
- 3) Does free volume theory, which has been very successful for predicting solvent and probe diffusion in non-glassy systems by relating the mobility of small molecules to the existence of void space, or "free volume", in solution, work as well for diffusion of probes below T_g ?

In order to answer these questions, we have measured D_{self} of two photochromic probes in polystyrene-toluene solutions over a wide range of temperature, above and below the glass transition. We have analyzed these data according to free volume theory for small molecule diffusion in non-glassy and glassy systems.

We have found, as expected, that the size of the diffusing molecule has a strong effect on its diffusion coefficient, and also that the diffusion coefficient of the probe in a polystyrene-toluene solution can be correlated to the toluene self-diffusion coefficient.

The effect of the glass transition on D_{self} has been measured. We have diffusion data from three PS toluene solutions, 81%, 83%, and 84% PS, versus temperature. There is clearly an abrupt change in the temperature dependence of D_{self} near T_g for each solution, with the abruptness of the change increasing with polymer concentration.

The predictions of the Vrentas-Duda free volume theory for glassy solutions have been examined. Although the theory is qualitatively correct in predicting the change in $D_{\text{self}}(T)$ at T_g and the effect of increasing polymer concentration on that change, the figure clearly shows that it does not agree quantitatively with the diffusion data. This is due to an inherently unrealistic representation of the polymer contribution to the glassy solution free volume. Thus, while the Vrentas-Duda theory is able to rationalize the diffusion data which we obtained, it is insufficient for a quantitative prediction of glassy self diffusion.

The substance of the work is described in two accompanying research reports.



Availability Codes	
Dist	Avail and/or Special
A-1	